



6352 CORTE DEL ABETO, STE B  
CARLSBAD, CA 92011 USA  
01.858.642.1988 • WWW.ATSBIO.COM

**Antibody to Human E-Selectin (CD62E)**  
MOUSE MONOCLONAL

**Catalog Number:** AB-438  
**Quantity:** 500 micrograms, 1 milligram  
**Format:** Lyophilized  
**Host:** Mouse  
**Isotype:** IgG1  
**Clone:** hCD62E  
**Immunogen:** Purified, Activated T-Cells

**Background:**

E-Selectin belongs to a family of divalent cation-dependent carbohydrate-binding glycoproteins or adhesion molecules. CD62E is expressed on the surface of endothelial cells and mediates the interaction of leukocytes and platelets with endothelial cells during an inflammatory response. CD62E is found in cytokine-stimulated endothelial cells and is accountable for the accumulation of blood leukocytes at sites of inflammation by mediating the adhesion of cells to the vascular lining. It demonstrates structural features such as the existence of lectin- and EGF-like domains followed by short consensus repeat (SCR) domains that contain 6 conserved cysteine residues. E-Selectin participates in the interaction between leukocytes and the endothelium and appears to be involved in the pathogenesis of atherosclerosis.

**Specificity and Preparation:**

Mouse anti-human E-Selectin (CD62E) is purified over an ion exchange column. Protein concentration is 1 mg/ml in PBS (after reconstitution).

**Usage and Storage:**

Reported to be effective as a staining antibody ( $10\mu\text{l}/10^6$  cells). Reconstitute with sterile H<sub>2</sub>O. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate. In lyophilized form, for long periods, store at 4°C. After reconstitution, if not intended for use within a month, aliquot and store at -20°C. Antibody is stable for two years lyophilized, one month in solution at 4°C.

**To view protocol(s) for this and other products please visit: [www.ATSBio.com/support/protocols](http://www.ATSBio.com/support/protocols)**