



## Protein G Recombinant

Catalog Number: PRP-647

**Quantity:** 1 milligram, 10 milligrams, 100 milligrams **Format:** Lyophilized powder containing no additives

**Host:** E. coli

## **Background:**

Protein G binds to the immunoglobulin constant region of many species. It can be used to detect, quantify and purify IgG antibodies and antibody/antigen complexes. Recombinant protein G contains only IgG binding domains. The albumin-binding domain as well as cell wall and cell membrane binding domains have been removed to ensure the maximum specific IgG binding capacity.

## **Specificity and Preparation:**

The recombinant protein G is produced in *Escherichia coli* using the sequence from Streptococcus C1-C2-C3. Recombinant protein G contains amino acids 190-384 of the native sequence, and has a molecular mass of 21.6 kDa. Recombinant protein-G migrates on SDS-PAGE around 32 kDa. It has a higher affinity for most mammalian immunoglobulins than protein A, including human IgG3 and rat IgG2a. It does not bind to human IgM, IgD, or IgA. Purity is >95% as determined by SDS-PAGE and RP-HPLC. Amino acid sequence:

MTYKLILNGKTLKGETTTEAVDAATAEKVFKQYANDNGVDGEWTYDDAT KTFTVTEKPEVIDASELTPAVTTYKLVINGKTLKGETTTEAVDAATAEKVFK QYANDNGVDGEWTYDDATKTFTVTEKPEVIDASELTPAVTTYKLVINGKTL KGETTTKAVDAETAEKAFKQYANDNGVDGVWTYDDATKTFTVTE.

## **Usage and Storage:**

Protein G is reported to be an excellent tool for purification and detection of IgG antibodies and antibody/antigen complexes.

Reconstitute with deionized water or PBS. After reconstitution, aliquot and store at -20°C. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

To view protocol(s) for this and other products please visit: www.ATSbio.com/support/protocols